

LEOPOLD KOCZAN and KATARZYNA TRĄBKA-WIECŁAW

Subclasses of typically real functions determined by some modular inequalities

ABSTRACT. Let T be the family of all typically real functions, i.e. functions that are analytic in the unit disk $\Delta := \{z \in \mathbb{C} : |z| < 1\}$, normalized by $f(0) = f'(0) - 1 = 0$ and such that $\operatorname{Im} z \operatorname{Im} f(z) \geq 0$ for $z \in \Delta$. Moreover, let us denote: $T^{(2)} := \{f \in T : f(z) = -f(-z) \text{ for } z \in \Delta\}$ and $T^{M,g} := \{f \in T : f \prec Mg \text{ in } \Delta\}$, where $M > 1$, $g \in T \cap S$ and S consists of all analytic functions, normalized and univalent in Δ .

We investigate classes in which the subordination is replaced with the majorization and the function g is typically real but does not necessarily univalent, i.e. classes $\{f \in T : f \ll Mg \text{ in } \Delta\}$, where $M > 1$, $g \in T$, which we denote by $T_{M,g}$. Furthermore, we broaden the class $T_{M,g}$ for the case $M \in (0, 1)$ in the following way: $T_{M,g} = \{f \in T : |f(z)| \geq M|g(z)| \text{ for } z \in \Delta\}$, $g \in T$.

1. Introduction. Let T be the family of all typically real functions, i.e. functions that are analytic in the unit disk $\Delta := \{z \in \mathbb{C} : |z| < 1\}$, normalized by $f(0) = f'(0) - 1 = 0$ and such that $\operatorname{Im} z \operatorname{Im} f(z) \geq 0$ for $z \in \Delta$. Let S denote the class of all analytic functions, normalized as above and univalent in Δ , and SR – the subclass of S consisting of functions with real coefficients. Moreover, let us denote: $T^{(2)} := \{f \in T : f(z) = -f(-z) \text{ for } z \in \Delta\}$ and $T^{M,g} := \{f \in T : f \prec Mg \text{ in } \Delta\}$, where $M > 1$, $g \in T \cap S$. The symbol $h \prec H$ denotes the subordination in Δ , i.e. $h(0) = H(0)$ and $h(\Delta) \subset H(\Delta)$, where H is univalent. Let us notice that for $g_1(z) = z$

2000 *Mathematics Subject Classification.* Primary 30C45. Secondary 30C80, 30C99.

Key words and phrases. Typically real functions, majorization, subordination.

and $g_2(z) = \frac{1}{2} \log \frac{1+z}{1-z}$ we have $\mathbf{T}^{M,g_1} = \{f \in \mathbf{T} : |f| < M \text{ in } \Delta\}$ and $\mathbf{T}^{M,g_2} = \{f \in \mathbf{T} : |\operatorname{Im} f| < M\pi/4 \text{ in } \Delta\}$, $M > 1$. These classes are briefly denoted by \mathbf{T}_M and $\mathbf{T}(M)$, respectively.

The subordination in the classes \mathbf{T} , \mathbf{S} and \mathbf{SR} has been investigated by several authors (for example [2], [3], [4]). The relation $\mathbf{T}^{M,g} = \{Mg(h/M) : h \in \mathbf{T}_M\}$ for $g \in \mathbf{T} \cap \mathbf{S}$ (see [3]) provides the following formula connecting different classes of type $\mathbf{T}^{M,g}$: $\mathbf{T}^{M,f} = \{Mf(g^{-1}(h/M)) : h \in \mathbf{T}^{M,g}\}$, $f, g \in \mathbf{T} \cap \mathbf{S}$. For this reason, instead of researching a class $\mathbf{T}^{M,f}$ one can consider a class $\mathbf{T}^{M,g}$, for instance \mathbf{T}_M or $\mathbf{T}(M)$. We apply this idea to obtain results in various classes $\mathbf{T}^{M,g}$ from corresponding results in the class $\mathbf{T}(M)$. Investigating $\mathbf{T}(M)$ is possible because the integral formula for this class, the set of extremal points and the set of supporting points are known (see [4]).

Moreover, it is easy to prove that the class $\mathbf{T}^{M,g} \cap \mathbf{T}^{(2)} = \{Mg(h/M) : h \in \mathbf{T}_M\}$ for $g \in \mathbf{T}^{(2)} \cap \mathbf{S}$.

In the paper we investigate classes similar to $\mathbf{T}^{M,g}$, in which the subordination is replaced with the majorization (the modular subordination) and the function g is typically real but does not necessarily univalent, i.e. classes $\mathbf{T}_{M,g} := \{f \in \mathbf{T} : f \ll Mg \text{ in } \Delta\}$, where $M > 1$, $g \in \mathbf{T}$. The symbol $h \ll H$ denotes the majorization in Δ , i.e. $|h(z)| \leq |H(z)|$ for all $z \in \Delta$.

Furthermore, we broaden the class $\mathbf{T}_{M,g}$ for the case when $M \in (0, 1)$ in the following way: $\mathbf{T}_{M,g} = \{f \in \mathbf{T} : |f(z)| \geq M|g(z)| \text{ for } z \in \Delta\}$, $g \in \mathbf{T}$.

Moreover, we study the subclass of the class $\mathbf{T}_{M,g}$, consisting of all odd functions, which we denote by $\mathbf{T}_{M,g}^{(2)}$.

The class $\mathbf{T}_{M,g}$ is not empty, because for example the function g belongs to this class. Analogously, the class $\mathbf{T}_{M,g}^{(2)}$ for $g \in \mathbf{T}^{(2)}$ is not empty. If $M = 1$, then the class consists of only one function g . So we investigate the class $\mathbf{T}_{M,g}$ for $M \in (0, 1) \cup (1, \infty)$. For $g = id$ and $M \geq 1$, we have $\mathbf{T}^{M,id} = \mathbf{T}_{M,id}$.

In the class $\mathbf{T}^{M,g}$ one can formulate theorems which are true for each function $g \in \mathbf{T} \cap \mathbf{S}$. However, in the class $\mathbf{T}_{M,g}$ it is impossible. Indeed, theorems in the class $\mathbf{T}_{M,g}$ in a fundamental way depends on the choice of the function g . It means that a theorem which is true in the class \mathbf{T}_{M,g_1} generally is not true in the class \mathbf{T}_{M,g_2} , for $g_1 \neq g_2$. In each case, we connect the researching class with the class \mathbf{T}_M or $\mathbf{T}_M^{(2)}$.

2. Some properties of the classes \mathbf{T} and $\mathbf{T}^{(2)}$. During our investigation of the class $\mathbf{T}_{M,g}$, we use the following relations of classes \mathbf{T} and $\mathbf{T}^{(2)}$, which we give as lemmas. In each lemma we shall prove only one implication. The other can be proved analogously. For simplicity, instead of h or $z \mapsto h(z)$ we will use $h(z)$.

Lemma 1. $f \in \mathbb{T} \iff \frac{1+z^2}{z}f(z^2) \in \mathbb{T}^{(2)}$.

Proof. Let $f \in \mathbb{T}$. For $f \in \mathbb{T}$ we have the Robertson formula $f(z) = \int_{-1}^1 \frac{z}{1-2zt+z^2} d\mu(t)$, where μ is a probability measure on $[-1, 1]$ (see [1], [2]). Then

$$\begin{aligned} \frac{(1+z^2)f(z^2)}{z} &= \int_{-1}^1 \frac{z(1+z^2)}{1-2z^2t+z^4} d\mu(t) = \int_{-1}^1 \frac{z(1+z^2)}{(1+z^2)^2-2(1+t)z^2} d\mu(t) \\ &= \int_0^1 \frac{z(1+z^2)}{(1+z^2)^2-4\tau z^2} d\nu(\tau) \end{aligned}$$

with $\nu(A) \equiv \mu(2A-1)$ (where A is a Borel set contained in $[0, 1]$). Clearly, $\int_0^1 \frac{z(1+z^2)}{(1+z^2)^2-4\tau z^2} d\nu(\tau) \in \mathbb{T}^{(2)}$ (the representation formula for functions from the class $\mathbb{T}^{(2)}$, see [5]). Therefore, $\frac{(1+z^2)f(z^2)}{z} \in \mathbb{T}^{(2)}$. \square

Lemma 2. $f \in \mathbb{T}^{(2)} \iff \frac{1+z^2}{1-z^2} \frac{f(iz)}{i} \in \mathbb{T}^{(2)}$.

Proof. Suppose that $f \in \mathbb{T}^{(2)}$. From Lemma 1, the function h given by $h(z^2) = \frac{z}{1+z^2} f(z)$ is in \mathbb{T} . The definition of h is correct since $h((-z)^2) = \frac{-z}{1+(-z)^2} f(-z) = \frac{zf(z)}{1+z^2} = h(z^2)$. Then $f(iz) = \frac{1-z^2}{iz} h(-z^2)$. Hence, $\frac{1+z^2}{1-z^2} \frac{f(iz)}{i} = -\frac{1+z^2}{z} h(-z^2)$. Because of Lemma 1 and the fact that $h \in \mathbb{T} \iff -h(-z) \in \mathbb{T}$, we receive $-\frac{1+z^2}{z} h(-z^2) \in \mathbb{T}^{(2)}$. This means that $\frac{1+z^2}{1-z^2} \frac{f(iz)}{i} \in \mathbb{T}^{(2)}$, so we have the desired result. \square

Lemma 3. $f \in \mathbb{T} \iff \frac{z^2}{(1-z^2)^2} \frac{1}{f(z)} \in \mathbb{T}$.

Proof. Let $f \in \mathbb{T}$. Then $f(z) = \frac{z}{1-z^2} p(z)$ for $p \in \text{PR}$ (the Rogosinski representation, [2], [6]), where PR consists of all analytic functions p such that $p(0) = 1$, $\text{Re } p(z) > 0$ for $z \in \Delta$ and having real coefficients. Clearly, $\frac{1}{p} \in \text{PR}$, so $\frac{z}{1-z^2} \frac{1}{p(z)} \in \mathbb{T}$, i.e. $\frac{z^2}{(1-z^2)^2} \frac{1}{f(z)} \in \mathbb{T}$. From this and the equality $\left\{ \frac{1}{p} : p \in \text{PR} \right\} = \text{PR}$, we get $f \in \mathbb{T} \iff \frac{z^2}{(1-z^2)^2} \frac{1}{f(z)} \in \mathbb{T}$. \square

Taking $f \in \mathbb{T}^{(2)}$ in Lemma 3, we obtain the following relation:

Lemma 4. $f \in \mathbb{T}^{(2)} \iff \frac{z^2}{(1-z^2)^2} \frac{1}{f(z)} \in \mathbb{T}^{(2)}$.

Lemma 5. $f \in \mathbb{T} \iff \frac{z^3}{(1-z^4)(1-z^2)} \frac{1}{f(z^2)} \in \mathbb{T}^{(2)}$.

Proof. Let $f \in \mathbb{T}$. On the basis of Lemma 1, the function g given by $g(z) = \frac{1+z^2}{z} f(z^2)$ belongs to $\mathbb{T}^{(2)}$. Hence, we have $\frac{z^2}{(1-z^2)^2} \frac{1}{g(z)} = \frac{z^3}{(1-z^4)(1-z^2)} \frac{1}{f(z^2)}$. From Lemma 4, we know that $\frac{z^2}{(1-z^2)^2} \frac{1}{g(z)} \in \mathbb{T}^{(2)}$ which is equivalent to $\frac{z^3}{(1-z^4)(1-z^2)} \frac{1}{f(z^2)} \in \mathbb{T}^{(2)}$. \square

Lemma 6. $f \in \mathbb{T}^{(2)} \iff \frac{z^2}{1-z^4} \frac{i}{f(iz)} \in \mathbb{T}^{(2)}$.

Proof. Suppose that $f \in \mathbb{T}^{(2)}$. Let $g(z) = \frac{1+z^2}{1-z^2} \frac{f(iz)}{i}$. By Lemma 2, $g \in \mathbb{T}^{(2)}$. Since $\frac{z^2}{(1-z^2)^2} \frac{1}{g(z)} = \frac{z^2}{1-z^4} \frac{i}{f(iz)}$, from Lemma 4 we get $\frac{z^2}{(1-z^2)^2} \frac{1}{g(z)} \in \mathbb{T}^{(2)}$ i.e. $\frac{z^2}{1-z^4} \frac{i}{f(iz)} \in \mathbb{T}^{(2)}$. \square

3. The majorization in the class of typically real functions \mathbb{T} . At the beginning we study the case when $M > 1$, i.e. the class

$$\mathbb{T}_{M,g} = \{f \in \mathbb{T} : |f(z)| \leq M|g(z)| \text{ for } z \in \Delta\}, \quad g \in \mathbb{T}.$$

At first, let $g(z) = \frac{z}{1+z}$. Clearly, $g \in \mathbb{T} \cap \mathbb{S}$.

Theorem 1. *If $f \in \mathbb{T}$ and $|f(z)| \leq M \left| \frac{z}{1+z} \right|$ for all $z \in \Delta$, $M > 1$ (i.e. $f \in \mathbb{T}_{M,g}$ where $g(z) = \frac{z}{1+z}$), then $f(z^2) \equiv \frac{z}{1+z^2} h(z)$ for some $h \in \mathbb{T}_M^{(2)}$.*

Proof. Let $f \in \mathbb{T}$ and $|f(z)| \leq M \left| \frac{z}{1+z} \right|$. Hence, $|f(z^2)| \leq M \left| \frac{z^2}{1+z^2} \right|$. Let $h(z) \equiv \frac{1+z^2}{z} f(z^2)$. By Lemma 1, $h \in \mathbb{T}^{(2)}$. Therefore, $f(z^2) \equiv \frac{z}{1+z^2} h(z)$. From the above equality, we get $\left| \frac{z}{1+z^2} \right| |h(z)| \leq M \left| \frac{z^2}{1+z^2} \right|$. This implies that $|h(z)| \leq M|z| < M$, that is $h \in \mathbb{T}_M^{(2)}$. \square

Now, let us consider the function $g(z) = z + z^3$. We have $g(z) = \frac{z}{1-z^2} (1 - z^4)$. Since $\operatorname{Re}(1 - z^4) > 0$ for $z \in \Delta$, from the Rogosinski formula (see [2], [6]), we get $g \in \mathbb{T}$. Moreover, $g \in \mathbb{T}^{(2)}$ and $g \notin \mathbb{S}$, because $g'(i/\sqrt{3}) = 0$.

Theorem 2. *If $f \in \mathbb{T}^{(2)}$ and $|f(z)| \leq M|z + z^3|$ for all $z \in \Delta$, $M > 1$ (i.e. $f \in \mathbb{T}_{M,g}^{(2)}$, where $g(z) = z + z^3$), then $f(z) \equiv \frac{1+z^2}{z} h(z^2)$ for some $h \in \mathbb{T}_M$.*

Proof. Suppose that $f \in \mathbb{T}^{(2)}$ and $|f(z)| \leq M|z + z^3|$. By Lemma 1, the function h given by $h(z^2) \equiv \frac{z}{1+z^2} f(z)$ is in \mathbb{T} . Therefore, $f(z) \equiv \frac{1+z^2}{z} h(z^2)$. From the second assumption, we have $\left| \frac{1+z^2}{z} \right| |h(z^2)| \leq M|z + z^3|$. Then $|h(z^2)| \leq M|z^2| < M$, i.e. $h \in \mathbb{T}_M$. \square

Let us study the next function $g(z) = \frac{z+z^3}{1-z^2}$. We have $g(z) = \frac{z}{1-z^2} (1+z^2)$. Since $\operatorname{Re}(1 + z^2) > 0$ for $z \in \Delta$, from the Rogosinski formula, $g \in \mathbb{T}$. Furthermore, $g \in \mathbb{T}^{(2)}$ and $g \notin \mathbb{S}$, because $g'(\sqrt{\sqrt{5}-2}i) = 0$.

Theorem 3. *If $f \in \mathbb{T}^{(2)}$ and $|f(z)| \leq M \left| \frac{z+z^3}{1-z^2} \right|$ for all $z \in \Delta$, $M > 1$ (i.e. $f \in \mathbb{T}_{M,g}^{(2)}$ where $g(z) = \frac{z+z^3}{1-z^2}$), then $f(z) \equiv \frac{1+z^2}{1-z^2} \frac{h(iz)}{i}$ for some $h \in \mathbb{T}_M^{(2)}$.*

Proof. Assume that $f \in \mathbb{T}^{(2)}$ and $|f(z)| \leq M \left| \frac{z+z^3}{1-z^2} \right|$. Let $h(iz) \equiv \frac{1-z^2}{1+z^2} if(z)$. By Lemma 2, $h \in \mathbb{T}^{(2)}$. Hence, $f(z) \equiv \frac{1+z^2}{1-z^2} \frac{h(iz)}{i}$. From the above equality, we get $\left| \frac{1+z^2}{1-z^2} \right| |h(iz)| \leq M \left| \frac{z+z^3}{1-z^2} \right|$. Therefore, $|h(iz)| \leq M|z| < M$, that is $h \in \mathbb{T}_M^{(2)}$. \square

In the further investigation we consider the case when $M \in (0, 1)$, i.e. the class

$$\mathbb{T}_{M,g} = \{f \in \mathbb{T} : |f(z)| \geq M|g(z)| \text{ for } z \in \Delta\}, \quad g \in \mathbb{T}.$$

Suppose that $g(z) = \frac{z}{(1-z^2)^2}$. Since $g(z) = \frac{z}{1-z^2} \frac{1}{1-z^2}$ and $\operatorname{Re} \left(\frac{1}{1-z^2} \right) > 0$ for $z \in \Delta$, hence $g \in \mathbb{T}$. We have also $g'(i/\sqrt{3}) = 0$, and it follows that $g \notin \mathbb{S}$.

Theorem 4. *If $f \in \mathbb{T}$ and $|f(z)| \geq M \left| \frac{z}{(1-z^2)^2} \right|$ for all $z \in \Delta$, $M \in (0, 1)$ (i.e. $f \in \mathbb{T}_{M,g}$ where $g(z) = \frac{z}{(1-z^2)^2}$), then $f(z) \equiv \frac{z^2}{(1-z^2)^2} \frac{1}{h(z)}$ for some $h \in \mathbb{T}_{1/M}$.*

Proof. Let $f \in \mathbb{T}$ and $|f(z)| \geq M \left| \frac{z}{(1-z^2)^2} \right|$. By Lemma 3, the function h given by $h(z) \equiv \frac{z^2}{(1-z^2)^2} \frac{1}{f(z)}$ belongs to \mathbb{T} . So $f(z) \equiv \frac{z^2}{(1-z^2)^2} \frac{1}{h(z)}$. From the second assumption, we have $\left| \frac{z^2}{(1-z^2)^2} \right| \frac{1}{|h(z)|} \geq M \left| \frac{z}{(1-z^2)^2} \right|$ i.e. $|h(z)| \leq |z|/M < 1/M$. Hence, $h \in \mathbb{T}_{1/M}$ and the proof is complete. \square

Analogously, using Lemma 4, we prove the following theorem:

Theorem 5. *If $f \in \mathbb{T}^{(2)}$ and $|f(z)| \geq M \left| \frac{z}{(1-z^2)^2} \right|$ for all $z \in \Delta$, $M \in (0, 1)$ (i.e. $f \in \mathbb{T}_{M,g}^{(2)}$ where $g(z) = \frac{z}{(1-z^2)^2}$), then $f(z) \equiv \frac{z^2}{(1-z^2)^2} \frac{1}{h(z)}$ for some $h \in \mathbb{T}_{1/M}^{(2)}$.*

Now, let us consider the function $g(z) = \frac{z}{(1-z^2)(1-z)}$. Clearly, $g(z) = \frac{z}{1-z^2} \frac{1}{1-z}$ and $\operatorname{Re} \left(\frac{1}{1-z} \right) > 0$ for $z \in \Delta$, so $g \in \mathbb{T}$. We have also

$$g' \left((i\sqrt{7} - 1)/4 \right) = 0,$$

which means that $g \notin \mathbb{S}$.

Theorem 6. *If $f \in \mathbb{T}$ and $|f(z)| \geq M \left| \frac{z}{(1-z^2)(1-z)} \right|$ for all $z \in \Delta$, $M \in (0, 1)$ (i.e. $f \in \mathbb{T}_{M,g}$ where $g(z) = \frac{z}{(1-z^2)(1-z)}$), then $f(z) \equiv \frac{z^3}{(1-z^4)(1-z^2)} \frac{1}{h(z)}$ for some $h \in \mathbb{T}_{1/M}^{(2)}$.*

Proof. Suppose that $f \in \mathbb{T}$ and $|f(z)| \geq M \left| \frac{z}{(1-z^2)(1-z)} \right|$. By Lemma 5, the function $h(z) \equiv \frac{z^3}{(1-z^4)(1-z^2)} \frac{1}{f(z^2)}$ is in $\mathbb{T}^{(2)}$. Hence, $f(z^2) \equiv \frac{z^3}{(1-z^4)(1-z^2)} \frac{1}{h(z)}$. From the second assumption, we get $\left| \frac{z^3}{(1-z^4)(1-z^2)} \right| \frac{1}{|h(z)|} \geq M \left| \frac{z^2}{(1-z^4)(1-z^2)} \right|$, so $|h(z)| \leq |z|/M < 1/M$. This means that $h \in \mathbb{T}_{1/M}^{(2)}$, so we have the desired result. \square

Now let us study the function $g(z) = \frac{z}{1-z^4}$. Because $g(z) = \frac{z}{1-z^2} \frac{1}{1+z^2}$ and $\operatorname{Re} \left(\frac{1}{1+z^2} \right) > 0$ for $z \in \Delta$, so $g \in \mathbb{T}$. Moreover, $g \in \mathbb{T}^{(2)}$ and $g \notin \mathbb{S}$, because $g'((i+1)/\sqrt[4]{12}) = 0$.

Theorem 7. *If $f \in \mathbb{T}^{(2)}$ and $|f(z)| \geq M \left| \frac{z}{1-z^4} \right|$ for all $z \in \Delta$, $M \in (0, 1)$ (i.e. $f \in \mathbb{T}_{M,g}^{(2)}$ where $g(z) = \frac{z}{1-z^4}$), then $f(iz) \equiv \frac{z^2}{1-z^4} \frac{i}{h(z)}$ for some $h \in \mathbb{T}_{1/M}^{(2)}$.*

Proof. Let $f \in \mathbb{T}^{(2)}$ and $|f(z)| \geq M \left| \frac{z}{1-z^4} \right|$. By Lemma 6, the function $h(z) \equiv \frac{z^2}{1-z^4} \frac{i}{f(iz)}$ belongs to $\mathbb{T}^{(2)}$. So $f(iz) \equiv \frac{z^2}{1-z^4} \frac{i}{h(z)}$. From the second assumption, we have $\left| \frac{z^2}{1-z^4} \right| \frac{1}{|h(z)|} \geq M \left| \frac{iz}{1-z^4} \right|$ i.e. $|h(z)| \leq |z|/M < 1/M$. Therefore, $h \in \mathbb{T}_{1/M}^{(2)}$ and the proof is complete. \square

The converses to Theorems 1–7 are also true.

REFERENCES

- [1] Duren, P. L., *Univalent Functions*, Springer-Verlag, New York, 1983.
- [2] Goodman, A. W., *Univalent Functions*, Mariner Publ. Co., Tampa, 1983.
- [3] Koczan, L., *On classes generated by bounded functions*, Ann. Univ. Mariae Curie-Skłodowska Sect. A **52** (2) (1998), 95–101.
- [4] Koczan, L., Szapiel, W., *Extremal problems in some classes of measures (IV). Typically real functions*, Ann. Univ. Mariae Curie-Skłodowska Sect. A **43** (1989), 55–68.
- [5] Koczan, L., Zaprawa, P., *On typically real functions with n -fold symmetry*, Ann. Univ. Mariae Curie-Skłodowska Sect. A **52** (2) (1998), 103–112.
- [6] Rogosinski, W. W., *Über positive harmonische Entwicklungen und tipisch-reelle Potenzreihen*, Math. Z. **35** (1932), 93–121 (German).

Leopold Koczan
 Department of Applied Mathematics
 Lublin University of Technology
 ul. Nadbystrzycka 38D
 20-618 Lublin
 Poland

Katarzyna Trąbka-Więclaw
 Department of Applied Mathematics
 Lublin University of Technology
 ul. Nadbystrzycka 38D
 20-618 Lublin
 Poland
 e-mail: k.trabka@pollub.pl

Received July 2, 2009