

RABHA W. IBRAHIM, MASLINA DARUS and NIKOLA TUNESKI

## On subordination for classes of non-Bazilevič type

ABSTRACT. We give some subordination results for new classes of normalized analytic functions containing differential operator of non-Bazilevič type in the open unit disk. By using Jack's lemma, sufficient conditions for this type of operator are also discussed.

**1. Introduction and preliminaries.** Consider the functions  $F$  in the open disk  $U := \{z \in \mathbb{C} : |z| < 1\}$ , defined by

$$\begin{aligned}
 (1.1) \quad F(z) &= \frac{z^\alpha}{(1-z)^\alpha} = \sum_{n=0}^{\infty} \frac{(\alpha)_n}{n!} z^{n+\alpha} \\
 &= z^\alpha + \sum_{n=1}^{\infty} \frac{(\alpha)_n}{n!} z^{n+\alpha} \\
 &= z^\alpha + \sum_{n=2}^{\infty} \frac{(\alpha)_{n-1}}{(n-1)!} z^{n+\alpha-1}, \quad \alpha \geq 1.
 \end{aligned}$$

From (1.1), assuming  $\alpha$  to be a parameter with the values  $\alpha := \frac{n+m}{m}$ ,  $m \in \mathbb{N}$ , and having  $n = 0$  in the first term of the series, we can write  $F$  in the form

$$(1.2) \quad F(z) = z + \sum_{n=2}^{\infty} \frac{(\alpha)_{n-1}}{(n-1)!} z^{n+\alpha-1}.$$

2000 *Mathematics Subject Classification.* 34G10, 26A33, 30C45.

*Key words and phrases.* Fractional calculus, subordination, non-Bazilevič function, Jack's lemma.

By employing (1.2), we define classes of analytic functions of fractional power.

Let  $\mathcal{A}_\alpha^+$  be the class of all normalized analytic functions  $F$  in the open disk  $U$  of the form

$$F(z) = z + \sum_{n=2}^{\infty} a_{n,\alpha} z^{n+\alpha-1}, \quad \alpha \geq 1,$$

satisfying  $F(0) = 0$  and  $F'(0) = 1$ . Moreover, let  $\mathcal{A}_\alpha^-$  be the class of all normalized analytic functions  $F$  in the open disk  $U$  of the form

$$F(z) = z - \sum_{n=2}^{\infty} a_{n,\alpha} z^{n+\alpha-1}, \quad a_{n,\alpha} \geq 0; \quad n = 2, 3, \dots,$$

satisfying  $F(0) = 0$  and  $F'(0) = 1$ .

**Definition 1.1** (Subordination Principle). For two functions  $f$  and  $g$  analytic in  $U$ , we say that the function  $f$  is subordinate to  $g$  in  $U$  and write  $f(z) \prec g(z)$  ( $z \in U$ ), if there exists a Schwarz function  $w(z)$  analytic in  $U$  with  $w(0) = 0$ , and  $|w(z)| < 1$ , such that  $f(z) = g(w(z))$ ,  $z \in U$ . In particular, if the function  $g$  is univalent in  $U$ , the above subordination is equivalent to  $f(0) = g(0)$  and  $f(U) \subset g(U)$ .

Now we define a differential operator as follows:

$$\begin{aligned} D_\alpha^0 F(z) &= F(z) = z + \sum_{n=2}^{\infty} a_{n,\alpha} z^{n+\alpha-1}, \quad \alpha \geq 1, \\ D_\alpha^1 F(z) &= \frac{F(z)}{2} + \frac{zF'(z)}{2} = z + \sum_{n=2}^{\infty} \frac{(n+\alpha)}{2} a_{n,\alpha} z^{n+\alpha-1}, \\ &\vdots \\ D_\alpha^k F(z) &= D(D^{k-1}F(z)) = z + \sum_{n=2}^{\infty} \left[ \frac{(n+\alpha)}{2} \right]^k a_{n,\alpha} z^{n+\alpha-1}. \end{aligned} \tag{1.3}$$

Let  $\mathcal{A}$  be the class of analytic functions of the form  $f(z) = z + a_2 z^2 + \dots$ . Obradović [8] introduced a class of functions  $f \in \mathcal{A}$  such that for  $0 < \mu < 1$ ,

$$\Re \left\{ f'(z) \left( \frac{z}{f(z)} \right)^\mu \right\} > 0, \quad z \in U. \tag{1.4}$$

He called it the class of function of non-Bazilevič type. There are many subordination results for this class (see [15]). In fact, this type of functions has been used to solve various problems (see [14]).

The main object of the present work is to apply a method based on the differential subordination in order to derive sufficient conditions for functions  $F \in \mathcal{A}_\alpha^+$  and  $F \in \mathcal{A}_\alpha^-$  to satisfy

$$(1.5) \quad (D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec q(z), \quad D_\alpha^k F(z) \neq 0, \quad z \in U,$$

where  $q$  is a given univalent function in  $U$  such that  $q(z) \neq 0$ ,  $\mu \neq 0$ .

Moreover, we give applications of these results in fractional calculus. We shall need the following known results:

**Lemma 1.1** ([4]). *Let  $q(z)$  be univalent in the unit disk  $U$  and  $\theta$  and  $\phi$  be analytic in a domain  $D$  containing  $q(U)$  with  $\phi(w) \neq 0$  when  $w \in q(U)$ . Set  $Q(z) := zq'(z)\phi(q(z))$ ,  $h(z) := \theta(q(z)) + Q(z)$ . Suppose that*

1.  $Q(z)$  is starlike univalent in  $U$ , and
2.  $\Re \frac{zh'(z)}{Q(z)} > 0$  for  $z \in U$ .

*If  $\theta(p(z)) + zp'(z)\phi(p(z)) \prec \theta(q(z)) + zq'(z)\phi(q(z))$ , then  $p(z) \prec q(z)$  and  $q$  is the best dominant.*

**Lemma 1.2** ([5]). *Let  $q(z)$  be convex univalent in the unit disk  $U$  and  $\psi$  and  $\gamma \in \mathbb{C}$  with  $\Re \left\{ 1 + \frac{zq''(z)}{q'(z)} + \frac{\psi}{\gamma} \right\} > 0$ . If  $p(z)$  is analytic in  $U$  and  $\psi p(z) + \gamma zp'(z) \prec \psi q(z) + \gamma zq'(z)$ , then  $p(z) \prec q(z)$  and  $q$  is the best dominant.*

**2. Subordination results.** In this section, we study subordination for normalized analytic functions in the classes  $\mathcal{A}_\alpha^+$  and  $\mathcal{A}_\alpha^-$ .

**Theorem 2.1.** *Let a function  $q$  be univalent in the unit disk  $U$  such that  $q(z) \neq 0$ ,  $\frac{zq'(z)}{q(z)}$  is starlike univalent in  $U$  and*

$$(2.1) \quad \Re \left\{ 1 + \frac{zq''(z)}{q'(z)} - \frac{zq'(z)}{q(z)} + \frac{a}{bq(z)} \right\} > 0, \quad b \neq 0, \quad q'(z) \neq 0, \quad z \in U.$$

*If  $F \in \mathcal{A}_\alpha^+$  satisfies the subordination*

$$\begin{aligned} \frac{a}{(D_\alpha^k F(z))'} \left( \frac{D_\alpha^k F(z)}{z} \right)^\mu + b \left[ \mu \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] \\ \prec \frac{a}{q(z)} + b \frac{zq'(z)}{q(z)}, \end{aligned}$$

*then*

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec q(z)$$

*and  $q$  is the best dominant.*

**Proof.** Let the function  $p$  be defined by

$$p(z) := (D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu, \quad D_\alpha^k F(z) \neq 0, \quad z \in U.$$

By setting

$$\theta(\omega) := \frac{a}{\omega} \text{ and } \phi(\omega) := \frac{b}{\omega}, \quad b \neq 0,$$

it can easily be observed that  $\theta(\omega)$  is analytic in  $\mathbb{C} - \{0\}$ ,  $\phi(\omega)$  is analytic in  $\mathbb{C} - \{0\}$  and that  $\phi(\omega) \neq 0$ ,  $\omega \in \mathbb{C} - \{0\}$ . Also we obtain

$$Q(z) = zq'(z)\phi(q(z)) = \frac{bzq'(z)}{q(z)}$$

and

$$h(z) = \theta(q(z)) + Q(z) = \frac{a}{q(z)} + b\frac{zq'(z)}{q(z)}.$$

It is clear that  $Q(z)$  is starlike univalent in  $U$ ,

$$\Re \left\{ \frac{zh'(z)}{Q(z)} \right\} = \Re \left\{ 1 + \frac{zq''(z)}{q'(z)} - \frac{zq'(z)}{q(z)} + \frac{a}{bq(z)} \right\} > 0.$$

By straightforward computation, we have

$$\begin{aligned} \frac{a}{p(z)} + b\frac{zp'(z)}{p(z)} &= \frac{a}{(D_\alpha^k F(z))'} \left( \frac{D_\alpha^k F(z)}{z} \right)^\mu \\ &\quad + b \left[ \mu \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] \\ &< \frac{a}{q(z)} + b\frac{zq'(z)}{q(z)}. \end{aligned}$$

Then by the assumption of the theorem, we see that the assertion of the theorem follows by application of Lemma 1.1.  $\square$

**Corollary 2.1.** *Assume that (2.1) holds and  $q$  is convex univalent in  $U$ . If  $F \in \mathcal{A}_\alpha^+$  and*

$$\begin{aligned} \frac{a}{(D_\alpha^k F(z))'} \left( \frac{D_\alpha^k F(z)}{z} \right)^\mu + b \left[ \mu \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] \\ < a \left( \frac{1+Bz}{1+Az} \right)^\mu + b \frac{\mu z(A-B)}{(1+Az)(1+Bz)}, \end{aligned}$$

then

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu < \left( \frac{1+Az}{1+Bz} \right)^\mu, \quad -1 \leq B < A \leq 1$$

and  $q(z) = \left( \frac{1+Az}{1+Bz} \right)^\mu$  is the best dominant.

**Corollary 2.2.** Assume that (2.1) holds and  $q$  is convex univalent in  $U$ . If  $F \in \mathcal{A}_\alpha^+$  and

$$\begin{aligned} \frac{a}{(D_\alpha^k F(z))'} \left( \frac{D_\alpha^k F(z)}{z} \right)^\mu + b \left[ \mu \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] \\ \prec a \left( \frac{1-z}{1+z} \right)^\mu + \frac{2\mu bz}{1-z^2}, \end{aligned}$$

for  $z \in U$ ,  $\mu \neq 0$ , then

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec \left( \frac{1+z}{1-z} \right)^\mu$$

and  $q(z) = \left( \frac{1+z}{1-z} \right)^\mu$  is the best dominant.

**Corollary 2.3.** Assume that (2.1) holds and  $q$  is convex univalent in  $U$ . If  $F \in \mathcal{A}_\alpha^+$  and

$$\begin{aligned} \frac{a}{(D_\alpha^k F(z))'} \left( \frac{D_\alpha^k F(z)}{z} \right)^\mu + b \left[ \mu \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] \\ \prec ae^{-\mu Az} + \mu bAz \end{aligned}$$

for  $z \in U$ ,  $\mu \neq 0$ , then

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec e^{\mu Az}$$

and  $q(z) = e^{\mu Az}$  is the best dominant.

The next result can be found in [3].

**Corollary 2.4.** Assume that  $k = 0$  in Theorem 2.1. Then

$$(F(z))' \left( \frac{z}{F(z)} \right)^\mu \prec q(z)$$

and  $q$  is the best dominant.

**Theorem 2.2.** Let a function  $q(z)$  be convex univalent in the unit disk  $U$  such that  $q'(z) \neq 0$  and

$$(2.2) \quad \Re \left\{ 1 + \frac{zq''(z)}{q'(z)} + \frac{1}{\gamma} \right\} > 0, \quad \gamma \neq 0.$$

Suppose that  $(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu$  is analytic in  $U$ . If  $F \in \mathcal{A}_\alpha^-$  satisfies the subordination

$$\begin{aligned} (D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \left[ \mu\gamma \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] \\ \prec q(z) + \gamma zq'(z), \end{aligned}$$

then

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec q(z), \quad z \in U, \quad D_\alpha^k F(z) \neq 0$$

and  $q$  is the best dominant.

**Proof.** Let the function  $p$  be defined by

$$p(z) := \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu, \quad D_\alpha^k F(z) \neq 0, \quad z \in U.$$

By setting  $\psi = 1$ , it can easily be observed that

$$\begin{aligned} & p(z) + \gamma z p'(z) \\ &= (D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \left[ \mu\gamma \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''(z)}{(D_\alpha^k F(z))'} \right] \\ &\prec q(z) + \gamma z q'(z). \end{aligned}$$

Then by the assumption of the theorem we see that the assertion of the theorem follows by application of Lemma 1.2.  $\square$

**Corollary 2.5.** Assume that (2.2) holds and  $q$  is convex univalent in  $U$ . If  $F \in \mathcal{A}_\alpha^-$  and

$$\begin{aligned} & (D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \left[ \mu\gamma \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''(z)}{(D_\alpha^k F(z))'} \right] \\ &\prec \left( \frac{1 + Az}{1 + Bz} \right)^\mu + \mu\gamma z(A - B) \frac{(1 + Az)^{\mu-1}}{(1 + Bz)^{\mu+1}}, \end{aligned}$$

then

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec \left( \frac{1 + Az}{1 + Bz} \right)^\mu, \quad -1 \leq B < A \leq 1$$

and  $q(z) = \left( \frac{1 + Az}{1 + Bz} \right)^\mu$  is the best dominant.

**Corollary 2.6.** Assume that (2.2) holds and  $q$  is convex univalent in  $U$ . If  $F \in \mathcal{A}_\alpha^-$  and

$$\begin{aligned} & (D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \left[ \mu\gamma \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''(z)}{(D_\alpha^k F(z))'} \right] \\ &\prec \left[ \frac{1 + z}{1 - z} \right]^\mu \left\{ 1 + \frac{2\gamma\mu z}{1 - z^2} \right\} \end{aligned}$$

for  $z \in U$ ,  $\mu \neq 0$ , then

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec \left( \frac{1 + z}{1 - z} \right)^\mu$$

and  $q(z) = \left( \frac{1 + z}{1 - z} \right)^\mu$  is the best dominant.

**Corollary 2.7.** *Assume that (2.2) holds and  $q$  is convex univalent in  $U$ . If  $F \in \mathcal{A}_\alpha^-$  and*

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \left[ \mu\gamma \left( 1 - \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} \right) + \frac{z(D_\alpha^k F(z))''(z)}{(D_\alpha^k F(z))'} \right] \prec e^{\mu Az} (1 + \mu\gamma Az)$$

for  $z \in U$ ,  $\mu \neq 0$ , then

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec e^{\mu Az}$$

and  $q(z) = e^{\mu Az}$  is the best dominant.

The next result can be found in [3].

**Corollary 2.8.** *Assume that  $k = 0$  in Theorem 2.2. Then*

$$(F(z))' \left( \frac{z}{F(z)} \right)^\mu \prec q(z)$$

and  $q$  is the best dominant.

**3. Applications.** In this section, we present some applications of Section 2 to fractional integral operators. Assume that  $f(z) = \sum_{n=2}^\infty \varphi_n z^{n-1}$  and let us begin with the following definitions:

**Definition 3.1** ([12]). The fractional integral of order  $\alpha$  is defined, for a function  $f$ , by

$$I_z^\alpha f(z) := \frac{1}{\Gamma(\alpha)} \int_0^z f(\zeta)(z - \zeta)^{\alpha-1} d\zeta, \quad \alpha \geq 1,$$

where the function  $f$  is analytic in a simply-connected region of the complex  $z$ -plane ( $\mathbb{C}$ ) containing the origin and the multiplicity of  $(z - \zeta)^{\alpha-1}$  is removed by requiring  $\log(z - \zeta)$  to be real when  $(z - \zeta) > 0$ .

Note that (see [12], [7])

$$I_z^\alpha z^\mu = \frac{\Gamma(\mu + 1)}{\Gamma(\mu + \alpha + 1)} z^{\mu+\alpha}, \quad (\mu > -1).$$

Thus we have

$$I_z^\alpha f(z) = \sum_{n=2}^\infty a_n z^{n+\alpha-1}$$

where  $a_n := \frac{\varphi_n \Gamma(n)}{\Gamma(n+\alpha)}$ , for all  $n = 2, 3, \dots$ . This implies that  $z + I_z^\alpha f(z) \in \mathcal{A}_\alpha^+$  and  $z - I_z^\alpha f(z) \in \mathcal{A}_\alpha^-$  ( $\varphi_n \geq 0$ ), so we get the following results:

**Theorem 3.1.** *Let the assumptions of Theorem 2.1 be satisfied. Then*

$$D_{\alpha}^k(z + I_z^{\alpha} f(z))' \left( \frac{z}{D_{\alpha}^k(z + I_z^{\alpha} f(z))} \right)^{\mu} \prec q(z), \quad z \neq 0, \quad z \in U$$

and  $q$  is the best dominant.

**Proof.** Consider the function  $F$  be defined by

$$F(z) := z + I_z^{\alpha} f(z), \quad z \in U, \quad z \neq 0. \quad \square$$

**Theorem 3.2.** *Let  $k = 0$  in Theorem 2.2. Then*

$$D_{\alpha}^k(z - I_z^{\alpha} f(z))' \left( \frac{z}{D_{\alpha}^k(z - I_z^{\alpha} f(z))} \right)^{\mu} \prec q(z), \quad z \neq 0, \quad z \in U$$

and  $q$  is the best dominant.

**Proof.** Consider the function  $F$  be defined by

$$F(z) := z - I_z^{\alpha} f(z), \quad z \in U, \quad z \neq 0. \quad \square$$

Let  $F(a, b; c; z)$  be the Gauss hypergeometric function (see [13]) defined, for  $z \in U$ , by

$$F(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n (1)_n} z^n,$$

where is the Pochhammer symbol defined by

$$(a)_n := \frac{\Gamma(a+n)}{\Gamma(a)} = \begin{cases} 1, & (n=0); \\ a(a+1)(a+2)\dots(a+n-1), & (n \in \mathbb{N}). \end{cases}$$

We need the following definition of fractional operators of the Saigo type fractional calculus (see [10], [9]).

**Definition 3.2.** For  $\alpha > 0$  and  $\beta, \eta \in \mathbb{R}$ , the fractional integral operator  $I_{0,z}^{\alpha, \beta, \eta}$  is defined by

$$I_{0,z}^{\alpha, \beta, \eta} f(z) = \frac{z^{-\alpha-\beta}}{\Gamma(\alpha)} \int_0^z (z-\zeta)^{\alpha-1} F\left(\alpha+\beta, -\eta; \alpha; 1-\frac{\zeta}{z}\right) f(\zeta) d\zeta$$

where the function  $f(z)$  is analytic in a simply-connected region of the  $z$ -plane containing the origin, with the order

$$f(z) = O(|z|^{\epsilon})(z \rightarrow 0), \quad \epsilon > \max\{0, \beta - \eta\} - 1$$

and the multiplicity of  $(z-\zeta)^{\alpha-1}$  is removed by requiring  $\log(z-\zeta)$  to be real when  $z-\zeta > 0$ .



From Definition 3.2, with  $\beta < 0$ , we have

$$\begin{aligned} I_{0,z}^{\alpha,\beta,\eta} f(z) &= \frac{z^{-\alpha-\beta}}{\Gamma(\alpha)} \int_0^z (z-\zeta)^{\alpha-1} F\left(\alpha+\beta, -\eta; \alpha; 1-\frac{\zeta}{z}\right) f(\zeta) d\zeta \\ &= \sum_{n=0}^{\infty} \frac{(\alpha+\beta)_n (-\eta)_n}{(\alpha)_n (1)_n} \frac{z^{-\alpha-\beta}}{\Gamma(\alpha)} \int_0^z (z-\zeta)^{\alpha-1} \left(1-\frac{\zeta}{z}\right)^n f(\zeta) d\zeta \\ &:= \sum_{n=0}^{\infty} B_n \frac{z^{-\alpha-\beta-n}}{\Gamma(\alpha)} \int_0^z (z-\zeta)^{n+\alpha-1} f(\zeta) d\zeta \\ &= \sum_{n=0}^{\infty} B_n \frac{z^{-\beta-1}}{\Gamma(\alpha)} f(\zeta) \\ &:= \frac{\bar{B}}{\Gamma(\alpha)} \sum_{n=2}^{\infty} \varphi_n z^{n-\beta-1} \end{aligned}$$

where  $\bar{B} := \sum_{n=0}^{\infty} B_n$ . Denote  $a_n := \frac{\bar{B}\varphi_n}{\Gamma(\alpha)}$ ,  $\forall n = 2, 3, \dots$ , and let  $\alpha = -\beta$ . Thus  $z + I_{0,z}^{\alpha,\beta,\eta} f(z) \in \mathcal{A}_\alpha^+$  and  $z - I_{0,z}^{\alpha,\beta,\eta} f(z) \in \mathcal{A}_\alpha^-$  ( $\varphi_n \geq 0$ ), so we have the following results:

**Theorem 3.3.** *Assume that the hypotheses of Theorem 2.1 are satisfied. Then*

$$D_\alpha^k(z + I_{0,z}^{\alpha,\beta,\eta} f(z))' \left( \frac{z}{D_\alpha^k(z + I_{0,z}^{\alpha,\beta,\eta} f(z))} \right)^\mu \prec q(z), \quad z \neq 0, z \in U$$

and  $q$  is the best dominant.

**Proof.** Consider the function  $F$  defined by

$$F(z) := z + I_{0,z}^{\alpha,\beta,\eta} f(z), \quad z \in U, z \neq 0. \quad \square$$

**Theorem 3.4.** *Assume that the hypotheses of Theorem 2.2 are satisfied. Then*

$$D_\alpha^k(z - I_{0,z}^{\alpha,\beta,\eta} f(z))' \left( \frac{z}{D_\alpha^k(z - I_{0,z}^{\alpha,\beta,\eta} f(z))} \right)^\mu \prec q(z), \quad z \neq 0, z \in U$$

and  $q$  is the best dominant.

**Proof.** Consider the function  $F$  defined by

$$F(z) := z - I_{0,z}^{\alpha,\beta,\eta} f(z), \quad z \in U, z \neq 0. \quad \square$$

**Remark 3.1.** Note that the authors have recently studied and defined several other classes of analytic functions related to fractional power (see [2], [1], [4]).

**4. The class  $\mathcal{S}_\mu(\gamma)$ .** A function  $F(z) \in \mathcal{A}_\alpha^+$  is said to be in the class  $\mathcal{S}_\mu(\gamma)$  if it satisfies

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec \frac{1+z}{1-\gamma z}, \quad (z \in U, \gamma \neq 1).$$

To discuss our problem, we have to recall here the following lemma due to Jack [15].

**Lemma 4.1.** *Let  $w$  be analytic in  $U$  with  $w(0) = 0$ . If  $|w(z)|$  attains its maximum value on the circle  $|z| = r < 1$  at a point  $z_0$ , then*

$$z_0 w'(z_0) = k w(z_0),$$

where  $k$  is a real number and  $k \geq 1$ .

We get the following result:

**Theorem 4.1.** *If  $F \in \mathcal{A}_\alpha^+$  satisfies*

$$(4.1) \quad \Re \left[ \mu - \mu \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] < \frac{1+\gamma}{2(1-\gamma)}, \quad (z \in U)$$

for some  $0 < \gamma < 1$ ,  $0 < \mu < 1$ , then  $F(z) \in \mathcal{S}_\mu(\gamma)$ .

**Proof.** Let  $w$  be defined by

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu = \frac{1+w(z)}{1-\gamma w(z)}, \quad (1 \neq \gamma w(z)).$$

Then  $w(z)$  is analytic in  $U$  with  $w(0) = 0$ . It follows that

$$\begin{aligned} \Re \left[ \mu - \mu \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] &= \Re \left[ \frac{z(\gamma w'(z) + 1)}{(1-\gamma w(z))(1+w(z))} \right] \\ &< \frac{1+\gamma}{2(1-\gamma)}, \quad \gamma \neq 1. \end{aligned}$$

Now we proceed to prove that  $|w(z)| < 1$ . Suppose that there exists a point  $z_0 \in U$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1.$$

Then, using Lemma 4.1 and letting  $w(z_0) = e^{i\theta}$  and  $z_0 w'(z_0) = k e^{i\theta}$ ,  $k \geq 1$ , we obtain

$$\begin{aligned} \Re \left[ \mu - \mu \frac{z(D_\alpha^k F(z_0))'}{D_\alpha^k F(z_0)} + \frac{z_0(D_\alpha^k F(z_0))''}{(D_\alpha^k F(z_0))'} \right] &= \Re \left[ \frac{z_0(w'(z_0)\gamma + 1)}{(1-\gamma w(z_0))(1+w(z_0))} \right] \\ &= \Re \left[ \frac{k e^{i\theta} \gamma + 1}{(1-\gamma e^{i\theta})(1+e^{i\theta})} \right] \\ &= \frac{k(\gamma + 1)}{2(1-\gamma)} \geq \frac{1+\gamma}{2(1-\gamma)}, \end{aligned}$$

$0 < \gamma < 1$ . Thus we have

$$\Re \left[ \mu - \mu \frac{z(D_\alpha^k F(z))'}{D_\alpha^k F(z)} + \frac{z(D_\alpha^k F(z))''}{(D_\alpha^k F(z))'} \right] \geq \frac{1 + \gamma}{2(1 - \gamma)}, \quad (z \in U)$$

which contradicts the hypothesis (4.1). Therefore, we conclude that  $|w(z)| < 1$  for all  $z \in U$  that is

$$(D_\alpha^k F(z))' \left( \frac{z}{D_\alpha^k F(z)} \right)^\mu \prec \frac{1 + z}{1 - \gamma z}, \quad (z \in U, \gamma \neq 1).$$

This completes the proof of the theorem.  $\square$

**Acknowledgement.** This work is supported by UKM-ST-06-FRGS0107-2009 and the authors would like to thank the referee for informative remarks given to improve the content of the paper.

#### REFERENCES

- [1] Darus, M., Ibrahim, R. W., *Coefficient inequalities for a new class of univalent functions*, Lobachevskii J. Math. **29**(4) (2008), 221–229.
- [2] Ibrahim, R. W., Darus, M., *On subordination theorems for new classes of normalize analytic functions*, Appl. Math. Sci. (Ruse) **2**(56) (2008), 2785–2794.
- [3] Ibrahim, R. W., Darus, M., *Subordination for new classes of non-Bazilevič type*, UNRI-UKM Symposium, KE-4 (2008).
- [4] Ibrahim, R. W., Darus, M., *Differential subordination results for new classes of the family  $\mathcal{E}(\Phi, \Psi)$* , JIPAM. J. Ineq. Pure Appl. Math. **10**(1) (2009), Art. 8, 9 pp.
- [5] Jack, I. S., *Functions starlike and convex of order  $k$* , J. London Math. Soc. **3** (1971), 469–474 .
- [6] Miller, S. S., Mocanu, P. T., *Differential Subordinations. Theory and Applications*, Monographs and Textbooks in Pure and Applied Mathematics, 225, Marcel Dekker, Inc., New York, 2000.
- [7] Miller, K. S., Ross, B., *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley and Sons, Inc., New York, 1993.
- [8] Obradović, M., *A class of univalent functions*, Hokkaido Math. J. **27**(2) (1998), 329–335.
- [9] Raina, R. K., *On certain class of analytic functions and applications to fractional calculus operator*, Integral Transform. Spec. Funct. **5** (1997), 247–260.
- [10] Raina, R. K., Srivastava, H. M., *A certain subclass of analytic functions associated with operators of fractional calculus*, Comput. Math. Appl. **32** (1996), 13–19.
- [11] Shanmugam, T. N., Ravichangran, V. and Sivasubramanian, S., *Differential sandwich theorems for some subclasses of analytic functions*, Austral. J. Math. Anal. Appl. **3**(1) (2006), 1–11.
- [12] Srivastava, H. M., Owa, S. (Eds.), *Univalent Functions, Fractional Calculus, and Their Applications*, Halsted Press, John Wiley and Sons, New York, Chichester, Brisbane, Toronto, 1989.
- [13] Srivastava, H. M., Owa, S. (Eds.), *Current Topics in Analytic Function Theory*, World Scientific Publishing Company, Singapore, New Jersey, London, Hong Kong, 1992.
- [14] Tuneski, N., Darus, M., *Fekete-Szegő functional for non-Bazilevič functions*, Acta Math. Acad. Paedagog. Nyházi. (N.S.) **18**(2) (2002), 63–65.

- [15] Wang, Z., Gao, C. and Liao, M., *On certain generalized class of non-Bazilevič functions*, Acta Math. Acad. Paedagog. Nyházi. (N.S.) **21**(2) (2005), 147–154.

Rabha W. Ibrahim  
School of Mathematical Sciences  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
Bangi 43600, Selangor Darul Ehsan  
Malaysia  
e-mail: rabhaibrahim@yahoo.com

Maslina Darus  
School of Mathematical Sciences  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
Bangi 43600, Selangor Darul Ehsan  
Malaysia  
e-mail: maslina@ukm.my (corresponding author)

Nikola Tuneski  
Faculty of Mechanical Engineering  
Karpoš II b.b., 1000 Skopje  
Republic of Macedonia  
e-mail: nikolat@mf.edu.mk

Received February 8, 2010